

ESSENTIALLY DISJOINT FAMILIES, CONFLICT FREE COLORINGS AND SHELAH'S REVISED GCH

LAJOS SOUKUP

ABSTRACT. Using Shelah's revised GCH theorem we prove that if $\mu < \beth_\omega \leq \lambda$ are cardinals, then every μ -almost disjoint family $\mathcal{A} \subset [\lambda]^{\beth_\omega}$ is *essentially disjoint*, i.e. for each $A \in \mathcal{A}$ there is a set $F(A) \in [A]^{<|A|}$ such that the family $\{A \setminus F(A) : A \in \mathcal{A}\}$ is disjoint.

We also show that if $\mu \leq \kappa \leq \lambda$ are cardinals, $\kappa \geq \omega$, and

- every μ -almost disjoint family $\mathcal{A} \subset [\lambda]^\kappa$ is essentially disjoint, then
- every μ -almost disjoint family $\mathcal{B} \subset [\lambda]^{\geq \kappa}$ has a *conflict-free coloring with κ colors*, i.e. there is a coloring $f : \lambda \rightarrow \kappa$ such that for all $B \in \mathcal{B}$ there is a color $\xi < \kappa$ such that $|\{\beta \in B : f(\beta) = \xi\}| = 1$.

Putting together these results we obtain that if $\mu < \beth_\omega \leq \lambda$, then every μ -almost disjoint family $\mathcal{B} \subset [\lambda]^{\geq \beth_\omega}$ has a conflict-free coloring with $\beth_\omega$ colors.

To yield the above mentioned results we also need to prove a certain compactness theorem concerning singular cardinals.

1. INTRODUCTION

The investigation and comparison of properties of almost families of sets has a long history, see e.g. the ancient paper of Miller, [10], the classical works of Erdős, Galvin, Hajnal and Rothchild, [1, 3, 4, 2], or the contributions of the next generation, [6, 7, 8] and [13].

A set system $\mathcal{A}$ is *μ -almost disjoint* iff $|A \cap A'| < \mu$ for distinct $A, A' \in \mathcal{A}$.

We say that $\mathcal{A} \subset [X]^\kappa$ is *essentially disjoint* (**ED**, in short) iff for each $A \in \mathcal{A}$ there is a set $F(A) \in [A]^{<\kappa}$ such that the family $\{A \setminus F(A) : A \in \mathcal{A}\}$ is disjoint.

$\mathbf{M}(\lambda, \kappa, \mu) \rightarrow \mathbf{ED}$ denotes the statement that every μ -almost disjoint family $\mathcal{A} \subset [\lambda]^\kappa$ is **ED**.

In [7, Theorems 2 and 5] Komjáth proved the following results:

- (A) $\mathbf{M}(\lambda, \omega, r) \rightarrow \mathbf{ED}$ for $r < \omega$;
- (B) $\mathbf{M}(\lambda, \omega_2, \omega) \rightarrow \mathbf{ED}$ under GCH;
- (C) $\mathbf{M}(\lambda, \omega_1, \omega) \rightarrow \mathbf{ED}$ if GCH holds and $\lambda \leq \aleph_\omega$;
- (D) $\mathbf{M}(\lambda, \omega_1, \omega) \rightarrow \mathbf{ED}$ if $V=L$.

Using Shelah' Revised GCH theorem, in Section 3 we will prove

Date: July 24, 2012.

2010 Mathematics Subject Classification. 03E05.

Key words and phrases. revised GCH, almost disjoint, essentially disjoint, sparse, conflict-free coloring, conflict-free chromatic number, singular cardinal compactness.

Research supported by the Hungarian National Research Grant OTKA K 83726.

(E) $\mathbf{M}(\lambda, \beth_\omega, \mu) \rightarrow \mathbf{ED}$ for each $\mu < \beth_\omega$.

If f is a function and A is a set, we let

$$I_f(A) = \{\xi \in \text{ran}(f) : |A \cap f^{-1}\{\xi\}| = 1\}.$$

A function f is a *conflict free coloring* of a set system $\mathcal{A}$ iff $\text{dom}(f) = \bigcup \mathcal{A}$ and $I_f(A) \neq \emptyset$ for all $A \in \mathcal{A}$.

$$\chi_{\text{CF}}(\lambda, \kappa, \mu) \leq \rho$$

denotes the statement that every μ -almost disjoint family $\mathcal{A} \subset [\lambda]^\kappa$ has a conflict-free coloring with ρ colors.

In [5] we proved that

- (a) $\chi_{\text{CF}}(\lambda, \kappa, r) \leq \omega$ for $r < \omega \leq \kappa \leq \lambda$;
- (b) $\chi_{\text{CF}}(\lambda, \kappa, \omega) \leq \omega_2$ for $\omega_2 \leq \kappa \leq \lambda$ under GCH;
- (c) $\chi_{\text{CF}}(\lambda, \kappa, \omega) \leq \omega_1$ for $\omega_1 \leq \kappa \leq \lambda \leq \aleph_\omega$ under GCH;
- (d) $\chi_{\text{CF}}(\lambda, \kappa, \omega) \leq \omega_1$ for $\omega_1 \leq \kappa \leq \lambda$ if $\mathbf{V}=\mathbf{L}$.

In [8, Corollary 2] Komjáth improved (a) by showing

- (a*) $\chi_{\text{CF}}(\lambda, \geq \omega, r) \leq \omega$ for $r < \omega \leq \kappa$,

i.e every r -almost disjoint family $\mathcal{A} \subset [\lambda]^{\geq \omega}$, where $r \in \omega$, has a conflict free coloring with ω colors.

In this paper we will show

- (b*) $\chi_{\text{CF}}(\lambda, \geq \omega_2, \omega) \leq \omega_2$ under GCH;
- (c*) $\chi_{\text{CF}}(\lambda, \geq \omega_1, \omega) \leq \omega_1$ if GCH holds and $\lambda \leq \aleph_\omega$;
- (d*) $\chi_{\text{CF}}(\lambda, \geq \omega_1, \omega) \leq \omega_1$ if $\mathbf{V}=\mathbf{L}$;
- (e*) $\chi_{\text{CF}}(\lambda, \geq \beth_\omega, \nu) \leq \beth_\omega$ for each $\nu < \beth_\omega$.

We do not need to present four proofs, because the following implication holds between essential disjointness and conflict free colorings:

$$\mathbf{M}(\lambda, \kappa, \mu) \rightarrow \mathbf{ED} \text{ implies } \chi_{\text{CF}}(\lambda, \geq \kappa, \mu) \leq \kappa, \quad (1)$$

see Corollary 4.2.

To prove our results, in section 2 we formulate a theorem which is a poor relative of Shelah's Singular Cardinal Compactness Theorem ([11, Theorem 2.1]).

Our notation is standard, see e.g. [9].

If X is a set and κ is a cardinal, then write $[X]^{\geq \kappa} = \{A \subset X : |A| \geq \kappa\}$.

If $\langle X_\alpha : \alpha < \sigma \rangle$ is a sequence of sets, we will often write $X_{<\alpha} = \bigcup_{\beta < \alpha} X_\beta$, $X_{\leq \alpha} = \bigcup_{\beta \leq \alpha} X_\beta$, etc.

2. POOR MAN'S SINGULAR CARDINAL COMPACTNESS

Both the formulation and the proof of the following statement use the ideas of Shelah's Singular Cardinal Compactness theorem (see [11, Theorem 2.1]).

A set system $\mathcal{G}$ is χ -chain closed iff $\bigcup_{\alpha < \chi} G_\alpha \in \mathcal{G}$ for any $\subseteq$ -increasing sequence $\langle G_\alpha : \alpha < \chi \rangle \subset \mathcal{G}$.

Theorem 2.1. *Assume that λ is a singular cardinal and $\mathcal{G} \subset [\lambda]^{<\lambda}$. If for some cardinals $\chi \leq \mu < \lambda$,*

- (o) *the set $\mathcal{G} \cap [\lambda]^\nu$ is χ -chain closed and cofinal in $[\lambda]^\nu$ for each $\mu \leq \nu < \lambda$,*

then there is a continuous, increasing sequence $\langle G_\xi : \xi < \text{cf}(\lambda) \rangle \subset \mathcal{G}$ such that $\bigcup_{\xi < \text{cf}(\lambda)} G_\xi = \lambda$.

Proof of Theorem 2.1. For each $Y \in \lambda$ with $\mu \leq |Y| < \lambda$ by (b) we can pick $G(Y) \in \mathcal{G} \cap [\lambda]^{|Y|}$ with $G(Y) \supset Y$.

Let $\langle \lambda_\zeta : \zeta < \text{cf}(\lambda) \rangle$ be a strictly increasing, continuous and cofinal sequence of cardinals in λ with $\lambda_0 > \mu, \chi, \text{cf}(\lambda)$.

By transfinite induction on $n < \chi$ we will define sets

$$\langle Y_{\zeta,n} : \zeta < \text{cf}(\lambda), n < \chi \rangle \subset [\lambda]^{<\lambda}$$

such that

- (A) $|Y_{\zeta,n}| = \lambda_\zeta$,
- (B) the sequences $\langle Y_{\zeta,n} : \zeta < \text{cf}(\lambda) \rangle$ are increasing and continuous,
- (C) $G(\bigcup_{m < n} Y_{\zeta,m}) \subset Y_{\zeta,n}$,

as follows (see Figure 1).

Let $Y_{\zeta,0} = \lambda_\zeta$ for $\zeta < \text{cf}(\lambda)$.

Assume that the family

$$\langle Y_{\zeta,m} : \zeta < \text{cf}(\lambda), m < n \rangle \subset [\lambda]^{<\lambda}$$

is defined.

Let

$$B_{\zeta,n} = G\left(\bigcup_{m < n} Y_{\zeta,m}\right),$$

and write

$$B_{\zeta,n} = \{b_{\zeta,n}(i) : i < \lambda_\zeta\}$$

for $\zeta < \text{cf}(\lambda)$, and for $\xi < \text{cf}(\lambda)$ let

$$Y_{\xi,n} = \{b_{\zeta,n}(i) : \zeta < \text{cf}(\lambda), i < \lambda_\xi\}.$$

Since the sequence $\langle \lambda_\zeta : \zeta < \text{cf}(\lambda) \rangle$ was continuous, the sequence $\langle Y_{\zeta,n} : \zeta < \text{cf}(\lambda) \rangle$ is also continuous, so (B) holds.

(A) and (C) are clear from the construction because $\text{cf}(\lambda) < \lambda_0$.

Let

$$G_\xi = \bigcup_{n < \chi} Y_{\xi,n}.$$

for $\xi < \text{cf}(\lambda)$.

To check $G_\xi \in \mathcal{G}$ it is enough to observe that

$$G_\xi = \bigcup_{n < \chi} G(Y_{\xi,n}).$$

because the sequence $\langle G(Y_{\xi,n}) : n < \chi \rangle \subset \mathcal{G} \cap [\lambda]^{\lambda_\xi}$ is $\subseteq$ -increasing and $\mathcal{G} \cap [\lambda]^{\lambda_\xi}$ is χ -chain closed.

By (B) we have $Y_{\zeta,n} \subset Y_{\xi,n}$ for $\zeta < \xi < \text{cf}(\lambda)$, and so $G_\zeta \subset G_\xi$.

Since $\lambda_\zeta \subset G_\zeta$, we also have $\bigcup_{\xi < \text{cf}(\lambda)} G_\xi = \lambda$.

Finally assume that $\xi < \text{cf}(\lambda)$ is a limit ordinal. Then

$$G_\xi = \bigcup_{n < \chi} Y_{\xi,n} = \bigcup_{n < \chi} \left(\bigcup_{\zeta < \xi} Y_{\zeta,n} \right) = \bigcup_{\zeta < \xi} \left(\bigcup_{n < \chi} Y_{\zeta,n} \right) = \bigcup_{\zeta < \xi} G_\zeta,$$




FIGURE 1.

so the sequence $\langle G_\xi : \xi < \text{cf}(\lambda) \rangle$ is continuous, which was to be proved. $\square$

3. FAMILIES OF SETS OF SIZE $\beth_\omega$

Definition 3.1. If $\nu \leq \rho$ are cardinals, then write

$$\rho^{[\nu]} = \rho$$

iff there is a family $\mathcal{B} \subset [\rho]^{<\nu}$ of size ρ such that for all $u \in [\rho]^\nu$ there is $\mathcal{P} \in [\mathcal{B}]^{<\nu}$ such that $u \subset \cup \mathcal{P}$.

Shelah's Revised CGH Theorem ([12, Theorem 0.1]).

If $\rho \geq \beth_\omega$, then $\rho^{[\nu]} = \rho$ for each large enough regular cardinal $\nu < \beth_\omega$.

Lemma 3.2. Fix $\mu < \beth_\omega$. Then for each $\rho \geq \beth_\omega$ there is a regular $\nu(\rho) < \beth_\omega$ such that if $\mathcal{A} \subset [\rho]^{\nu(\rho)}$ is μ -almost disjoint, then $|\mathcal{A}| \leq \rho$.

Proof of lemma 3.2. Let $\mu < \nu < \beth_\omega$ be regular such that $\rho^{[\nu]} = \rho$ witnessed by a family $\mathcal{B} \subset [\rho]^\nu$. We show that $\nu(\rho) = \nu$ works.

If $A \in \mathcal{A}$, then there is $B \in \mathcal{B}$ such that $A \cap B = \nu$. If $A_0, A_1 \in \mathcal{A}$ are distinct, and $|A_0 \cap B| = |A_1 \cap B| = \nu$ then $A_0 \cap B \neq A_1 \cap B$ because $\mathcal{A}$ is μ -almost disjoint. So

$$|\{A \in \mathcal{A} : |A \cap B| = \nu\}| \leq 2^{|B|} \leq 2^\nu < \beth_\omega.$$

Thus $|\mathcal{A}| \leq \beth_\omega \cdot |\mathcal{B}| = \rho$. $\square$

Lemma 3.3. If $\lambda > \beth_\omega > \mu$, and $\{A_\alpha : \alpha < \tau\} \subset [\lambda]^{\beth_\omega}$ is a μ -almost disjoint family, then $\tau \leq \lambda$, and there is an increasing, continuous sequence $\langle G_\zeta : \zeta < \text{cf}(\lambda) \rangle \subset [\lambda]^{<\lambda}$ such that

$$\forall \zeta < \text{cf}(\lambda) \quad \forall \alpha \in G_{\zeta+1} \setminus G_\zeta \quad (|A_\alpha \cap G_\zeta| < \beth_\omega \text{ and } A_\alpha \subset G_{\zeta+1}). \quad (2)$$

Proof. Applying lemma 3.2 for $\rho = \lambda$ we have $|\tau| \leq \lambda$. So we can assume that $\tau = \lambda$.

We should distinguish two cases.

Case 1. λ is a regular cardinal

Denoting by **Card** the class of cardinals, pick a cardinal $\mu < \nu < \beth_\omega$ such that

$$B = \{\rho \in \mathbf{Card} \cap \lambda : \nu(\rho) = \nu\}$$

is cofinal in **Card** $\cap \lambda$. (If $\lambda = \sigma^+$ is a successor cardinal, then $B = \{\sigma\}$ and $\nu = \nu(\sigma)$ work.)

Let

$$B^* = \{\zeta < \lambda : |\zeta| \in B\},$$

and for $\zeta \in B^*$ let

$$f(\zeta) = A_{<\zeta} \cup \{\alpha : |A_\alpha \cap \zeta| \geq \nu\}.$$

Since the family $\{A_\alpha \cap \zeta : |A_\alpha \cap \zeta| \geq \nu\} \subset [\zeta]^{>\nu}$ is μ -almost disjoint, by Lemma 3.2 we have $|f(\zeta)| \leq |\zeta|$. Let

$$D = \{\delta < \lambda : \text{cf}(\delta) = \nu^+, \sup(B^* \cap \delta) = \delta \wedge (\forall \zeta \in B^* \cap \delta) f(\zeta) \subset \delta\}.$$

Since B^* is cofinal in λ , $\nu^+ \leq \beth_\omega < \lambda$ and $f : B^* \rightarrow [\lambda]^{<\lambda}$, the set D is also cofinal in λ , and

$$\forall \delta \in D (\forall \alpha < \delta A_\alpha \subset \delta \wedge \forall \alpha \in (\lambda \setminus \delta) |A_\alpha \cap \delta| < \nu).$$

Indeed, if $|A_\alpha \cap \delta| \geq \nu$, then there is $\zeta \in B^* \cap \delta$ with $|A_\alpha \cap \delta| \geq \nu$ for $\text{cf}(\delta) = \nu^+$, and then $\alpha \in f(\zeta) \subset \delta$ by the definition of f .

Thus we also have

$$\forall \gamma \in D' (\forall \alpha < \gamma A_\alpha \subset \gamma \wedge \forall \alpha \in (\lambda \setminus \gamma) |A_\alpha \cap \gamma| < \beth_\omega). \quad (3)$$

Indeed, if $|A_\alpha \cap \gamma| = \beth_\omega$, then there is $\delta \in D \cap (\gamma + 1)$ with $|A_\alpha \cap \delta| \geq \nu$, and so $A_\alpha \subset \delta$.

Let $\{\gamma_\zeta : \zeta < \lambda\}$ be the increasing enumeration of the club set D' . Then the choice $G_\zeta = \gamma_\zeta$ works by (3).

Case 2. $\lambda > \text{cf}(\lambda)$ is a singular cardinal.

Let

$$\mathcal{G} = \{G \in [\lambda]^{<\lambda} : |G| \geq \beth_\omega \wedge (\forall \alpha \in G) A_\alpha \subset G \wedge (\forall \alpha \in \lambda \setminus G) |A_\alpha \cap G| < \nu(|G|)\}.$$

If $\beth_\omega \leq \rho < \lambda$, then $\nu(\rho) > \omega$ is a regular cardinal, so the family $\mathcal{G} \cap [\lambda]^\rho$ is ω -chain closed.

Next we show that the set $\mathcal{G} \cap [\lambda]^\rho$ is cofinal in $[\lambda]^\rho$ for all $\beth_\omega \leq \rho < \lambda$.

Indeed, let $Y \in [\lambda]^\rho$. Define an increasing sequence $\langle Y_n : n < \omega \rangle \subset [\lambda]^\rho$ as follows:

- (i) $Y_0 = Y$,
- (ii) for $1 \leq n < \omega$ let

$$Y_n = Y_{<n} \cup \bigcup \{A_\alpha : \alpha \in Y_{n-1}\} \cup \{\alpha : |Y_{<n} \cap A_\alpha| \geq \nu(\rho)\}.$$

We are to show that $G = \bigcup_{n < \omega} Y_n \in \mathcal{G} \cap [\lambda]^\rho$.

By induction on n we obtain $|Y_n| = \rho$ because $|\{\alpha : |Y_{<n} \cap A_\alpha| \geq \nu(\rho)\}| \leq \rho$ by lemma 3.2. So $|G| = \rho$.

If $\alpha \in G$, then $\alpha \in Y_n$ for some n , and so $A_\alpha \subset Y_{n+1} \subset G$.

If $|A_\alpha \cap G| \geq \nu(\rho)$, then $|A_\alpha \cap Y_n| \geq \nu(\rho)$ for some $n < \omega$ because $\text{cf}(\nu(\rho)) = \nu(\rho) > \omega$. So $\alpha \in Y_{n+1} \subset G$.

Thus we proved $G \in \mathcal{G}$ and so $\mathcal{G} \cap [\lambda]^\rho$ is really cofinal in $[\lambda]^\rho$.

So we can apply Theorem 2.1 to obtain the required sequence $\langle G_\zeta : \zeta < \text{cf}(\lambda) \rangle$. $\square$

Theorem 3.4. $\mathbf{M}(\lambda, \beth_\omega, \mu) \rightarrow \mathbf{ED}$ whenever $\nu < \beth_\omega \leq \lambda$.

Proof. We prove the theorem by induction on λ . If $\lambda = \beth_\omega$, then applying lemma 3.2 for $\rho = \beth_\omega$ we have $|\mathcal{A}| \leq \beth_\omega$, so the statement is clear.

Assume now that $\lambda > \beth_\omega$ and we proved the theorem for $\lambda' < \lambda$.

Let $\mathcal{A} \subset [\lambda]^{\beth_\omega}$ be μ -almost disjoint. By lemma 3.2, $|\mathcal{A}| \leq \lambda$, so we can write $\mathcal{A} = \{A_\alpha : \alpha < \lambda\}$.

By lemma 3.3 there is an increasing, continuous sequence $\{G_\zeta : \zeta < \text{cf}(\lambda)\} \subset [\lambda]^{<\lambda}$ which satisfies the (2).

For $\zeta < \text{cf}(\lambda)$ write

$$X_\zeta = G_{\zeta+1} \setminus G_\zeta \text{ and } \mathcal{A}_\zeta = \{A_\alpha \cap X_\zeta : \alpha \in X_\zeta\}.$$

Then $\mathcal{A}_\zeta \subset [X_\zeta]^{\beth_\omega}$ is μ -almost disjoint, so by the inductive assumption there are set $F_\zeta(A_\alpha) \in [A_\alpha]^{<\beth_\omega}$ for $\alpha \in X_\zeta$ such that the family

$$\{(A_\alpha \cap X_\zeta) \setminus F_\zeta(A_\alpha) : \alpha \in X_\zeta\}$$

is disjoint. Let

$$F(A_\alpha) = F_\zeta(A_\alpha) \cup (A_\alpha \cap X_{<\zeta})$$

for $\alpha \in X_\zeta$. Then $|F(A_\alpha)| < \beth_\omega$, and the sets

$$\{A_\alpha \setminus F(A_\alpha) : \alpha < \lambda\}$$

are disjoint. $\square$

4. ESSENTIAL DISJOINTNESS AND CONFLICT FREE COLORINGS

We say that a set system $\mathcal{A}$ is κ -hereditary essentially disjoint, (κ -**hED**, in short) iff for each function H with $\text{dom}(H) = \mathcal{A}$ and $H(A) \in [A]^\kappa$ for $A \in \mathcal{A}$, the family $\{H(A) : A \in \mathcal{A}\}$ is **ED**.

A κ -**hED** family is clearly κ -almost disjoint.

Theorem 4.1. Let $\lambda \geq \kappa \geq \omega$ be cardinals and $\mathcal{A} \subset [\lambda]^{\geq \kappa}$ be a family of sets. If

- $\mathcal{A}$ is κ -hereditarily essentially disjoint,

then

- there is a coloring $c : \lambda \rightarrow \kappa$ such that $|I_c(A)| = \kappa$ for all $A \in \mathcal{A}$.

Corollary 4.2. Let $\mu \leq \kappa \leq \lambda$ be cardinals, $\kappa \geq \omega$. If

$(*_\lambda)$ every μ -almost disjoint family $\mathcal{A} \subset [\lambda]^\kappa$ is **ED**,

then

$(\star_\lambda)$ for each μ -almost disjoint family $\mathcal{A} \subset [\lambda]^{\geq \kappa}$ there is a coloring $f : \lambda \rightarrow \kappa$ such that $|I_f(A)| = \kappa$ for all $A \in \mathcal{A}$, and so $\chi_{\text{CF}}(\lambda, \geq \kappa, \mu) \leq \kappa$.

In the inductive proof of Theorem 4.1 we will use the following observation on decomposability of κ -**hED** families.

Theorem 4.3. *If $\lambda > \kappa \geq \omega$ are cardinals, and $\mathcal{A} = \{A_\alpha : \alpha < \lambda\} \subset [\lambda]^{\geq \kappa}$ is a κ -**hED** family, then there is a partition $\{X_\zeta : \zeta < \text{cf}(\lambda)\} \subset [\lambda]^{< \lambda}$ of λ such that*

- (M1) $A_\alpha \cap A_{\alpha'} \subset X_{\leq \zeta}$ for distinct $\alpha, \alpha' \in X_{\leq \zeta}$.
- (M2) $|A_\alpha \cap X_{< \zeta}| < \kappa$ for all $\alpha \in X_\zeta$,
- (M3) $|A_\alpha \cap X_\zeta| \geq \kappa$ for all $\alpha \in X_\zeta$,

Proof of Theorem 4.3. We should distinguish two cases.

Case 1. λ is regular.

For each $A \in \mathcal{A}$, let $H(A)$ be the first κ many elements of A . Since the family $\{H(A) : A \in \mathcal{A}\}$ is **ED**, there is an injective function f on $\mathcal{A}$ with $f(A) \in H(A)$.

Fix a large enough regular cardinal θ , and let $\langle N_\zeta : 1 \leq \zeta < \lambda \rangle$ be a strictly increasing continuous sequence of elementary submodels of $\langle H_\theta, \in \rangle$ such that

- (i) $\kappa + \zeta \subset N_\zeta \cap \lambda \in \lambda$ and $|N_\zeta| = \kappa + |\zeta|$,
- (ii) $\langle A_\alpha : \alpha < \lambda \rangle, H, f \in N_1$.
- (iii) $N_\zeta \in N_{\zeta+1}$.

Write $N_0 = \emptyset$.

For $\zeta < \text{cf}(\lambda)$ let

$$X_\zeta = (N_{\zeta+1} \setminus N_\zeta) \cap \lambda.$$

(M1) is clear because $N_{\zeta+1}$ is an elementary submodel, $X_{\leq \zeta} = N_{\zeta+1} \cap \lambda$, and $\kappa + 1 \subset N_{\zeta+1}$.

To check (M2), assume that $\alpha < \lambda$ with $|A_\alpha \cap X_{< \zeta}| = \kappa$. Since $H(A_\alpha)$ is the first κ many elements of A_α , and $X_{< \zeta} = N_\zeta \cap \lambda$ is an initial segment of λ , we have $H(A_\alpha) \subset N_\zeta$. So $\eta = f(A_\alpha) \in H(A_\alpha) \subset N_\zeta$. But α is definable from f , $\langle A_\alpha : \alpha < \lambda \rangle$ and η , because f was injective. So $\alpha \in N_\zeta \cap \lambda = X_{< \zeta}$.

To check (M3) assume that $\alpha \in (N_{\zeta+1} \setminus N_\zeta) \cap \lambda$. Then $|A_\alpha \cap N_{\zeta+1}| \geq \kappa$ because $N_{\zeta+1}$ is an elementary submodel and $\kappa + 1 \subset N_{\zeta+1}$. On the other hand, $|A \cap N_\zeta| < \kappa$ by (M2), so $|A_\alpha \cap (N_{\zeta+1} \setminus N_\zeta)| = \kappa$.

Case 2. λ is singular.

Let

$$\chi = \begin{cases} \omega & \text{cf}(\kappa) \neq \omega, \\ \omega_1 & \text{cf}(\kappa) = \omega. \end{cases}$$

We say that $Y \subset \lambda$ is *good* if (G1)-(G3) below hold:

- (G1) if $\alpha \neq \beta \in Y$, then $A_\alpha \cap A_\beta \subset Y$,
- (G2) if $\alpha \in Y$ then $|A_\alpha \cap Y| \geq \kappa$,
- (G3) if $|A_\alpha \cap Y| \geq \kappa$ then $\alpha \in Y$.

Let

$$\mathcal{G} = \{Y \in [\lambda]^{<\lambda} : Y \text{ is good}\}.$$

Lemma 4.4. $\mathcal{G} \cap [\lambda]^\nu$ is χ -chain closed for $\max(\kappa, \text{cf}(\lambda), \chi) < \nu < \lambda$.

Proof. Assume that $\langle Y_n : n < \chi \rangle \subset \mathcal{G} \cap [\lambda]^\nu$ is increasing.

(G1) and (G2) are clear.

To check (G3) assume that $|A_\alpha \cap \bigcup_{n < \chi} Y_n| \geq \kappa$. Since $\text{cf}(\kappa) \neq \chi$, there is $n < \chi$ with $|A_\alpha \cap Y_n| = \kappa$, and so $\alpha \in Y_n$. $\square$

Lemma 4.5. $\mathcal{G} \cap [\lambda]^\nu$ is cofinal in $[\lambda]^\nu$ for $\max(\kappa, \text{cf}(\lambda), \chi) < \nu < \lambda$.

Proof of the Lemma. Let $Y \in [\lambda]^\nu$.

For each $A \in \mathcal{A}$ pick $H(A) \in [A]^\kappa$.

Define an increasing sequence $\langle Y_n : n < \chi \rangle \subset [\lambda]^\nu$ as follows:

- (i) $Y_0 = Y$,
- (ii) for $1 \leq n < \chi$ let

$$Y_n = Y_{<n} \cup \bigcup \{A_\alpha \cap A_\beta : \alpha \neq \beta \in Y_{<n}\} \cup \bigcup \{H(A_\alpha) : \alpha \in Y_{<n}\} \cup \{\alpha : |Y_{<n} \cap A_\alpha| \geq \kappa\}.$$

Let $G = \bigcup_{n < \chi} Y_n$. We are to show that $G \in \mathcal{G} \cap [\lambda]^\nu$.

Since $|Y_n| \leq |Y_{<n}| + \kappa$ by inequality (5), we have $\nu \leq |Y| \leq |G| \leq |Y| + \kappa + \chi = \nu$.

Properties (G1) and (G2) are straightforward from the construction.

To check (G3) assume that $|A_\alpha \cap G| \geq \kappa$. Since $\text{cf}(\kappa) \neq \chi$, there is $n < \chi$ with $|A_\alpha \cap Y_n| = \kappa$, and so $\alpha \in Y_{n+1} \subset G$. $\square$

By Theorem 2.1 there is a continuous, increasing sequence $\langle G_\xi : \xi < \text{cf}(\lambda) \rangle \subset \mathcal{G}$ such that $\bigcup_{\xi < \text{cf}(\lambda)} G_\xi = \lambda$.

Now for $\zeta < \text{cf}(\lambda)$ let

$$X_\zeta = G_{\zeta+1} \setminus G_\zeta.$$

Clearly $|X_\zeta| \leq |G_\zeta| < \lambda$. The sequence $\langle G_\xi : \xi < \text{cf}(\lambda) \rangle$ was continuous, so the family $\{X_\zeta : \zeta < \text{cf}(\lambda)\}$ is a partition of λ .

Since $G_\zeta \in \mathcal{G}$ we have

$$|A_\alpha \cap G_\zeta| \geq \kappa \text{ iff } \alpha \in G_\zeta. \quad (4)$$

(M1) is clear because $X_{\leq \zeta} = G_{\zeta+1} \in \mathcal{G}$ and (G1) holds for $G_{\zeta+1}$.

To check (M2) assume that $\alpha \in X_\zeta$, and so $\alpha \notin G_\zeta$. Thus $|A_\alpha \cap G_\zeta| < \kappa$ by (4). But $G_\zeta = X_{<\zeta}$, so (M2) holds.

To check (M3) assume that $\alpha \in X_\zeta$. Then $\alpha \in G_{\zeta+1} \setminus G_\zeta$ and so applying (4) twice we obtain

$$|A_\alpha \cap G_{\zeta+1}| \geq \kappa \wedge |A_\alpha \cap G_\zeta| < \kappa.$$

So $|A_\alpha \cap X_\zeta| \geq \kappa$.

So we proved Theorem 4.3. $\square$

Proof of Theorem 4.1. Let $\mathcal{I}$ be a principal ideal on κ with $[\kappa]^{<\kappa} \subset \mathcal{I}$. We prove the following stronger statement:

- ($\bullet_\lambda$) If a family $\mathcal{A} \subset [\lambda]^{\geq \kappa}$ is κ -**hED**, and $F : \lambda \rightarrow \mathcal{I}$, then there is a function $c : \lambda \rightarrow \kappa$ such that
- ($\mathcal{I}1$) $c(\xi) \notin F(\xi)$ for $\xi \in \lambda$,
 - ($\mathcal{I}2$) $|I_c(A)| = \kappa$ for all $A \in \mathcal{A}$.

If $Y \subset \lambda$, then consider the family

$$\mathcal{A}[Y] = \{A \in \mathcal{A} : |A \cap Y| \geq \kappa\},$$

and for each $A \in \mathcal{A}[Y]$ pick $H(A) \in [A \cap Y]^\kappa$. Since $\mathcal{A}$ is κ -**hED**, the family $\{H(A) : A \in \mathcal{A}[Y]\} \subset [Y]^\kappa$ is essentially disjoint, so

$$|\mathcal{A}[Y]| \leq |Y|. \quad (5)$$

Especially, $|\mathcal{A}| \leq \lambda$, and so we can write $\mathcal{A} = \{A_\alpha : \alpha < \lambda\}$.

We prove ($\bullet_\lambda$) by induction on λ .

If $\lambda = \kappa$, then let $c : \lambda \rightarrow \kappa$ be any injective function such that $c(\xi) \notin F(\xi)$ for $\xi \in \lambda$. Then $I_c(A) = c''A \in [\kappa]^\kappa$ for all $A \in \mathcal{A}$.

Assume now that $\lambda > \kappa$ and ($\bullet_{\lambda'}$) holds for $\lambda' < \lambda$.

Apply Theorem 4.3 for the family $\{A_\alpha : \alpha < \lambda\}$ to find a partition $\{X_\zeta : \zeta < \text{cf}(\lambda)\} \subset [\lambda]^{<\lambda}$ of λ satisfying (M1)–(M3).

Let $\mathcal{A}_\zeta = \{A_\alpha : \alpha \in X_\zeta\} \in [\mathcal{A}]^{<\lambda}$ for $\zeta < \text{cf}(\lambda)$.

By recursion, for each $\zeta < \text{cf}(\lambda)$ we define a function

$$c_\zeta : X_\zeta \rightarrow \kappa$$

and a function $G_\zeta : \mathcal{A}_\zeta \rightarrow \mathcal{I} \cap [\kappa]^\kappa$ such that

$$G_\zeta(A_\alpha) \subset I_{c_\zeta}(A_\alpha) \text{ for all } \alpha \in X_\zeta \quad (6)$$

as follows.

Assume that we have defined $\langle c_\eta : \eta < \zeta \rangle$ and $\langle G_\eta : \eta < \zeta \rangle$.

If $x \in X_\zeta$ then there is at most one $\alpha \in X_{<\zeta}$ with $x \in A_\alpha$ by (M1). Define $F_\zeta : X_\zeta \rightarrow \mathcal{I}$ as follows:

$$F_\zeta(x) = \begin{cases} F(x) \cup G_\eta(A_\alpha) & \text{if } x \in A_\alpha \in \mathcal{A}_\eta \text{ for some } \eta < \zeta, \\ F(x) & \text{otherwise.} \end{cases}$$

Then, by the inductive assumption ($\bullet_{|X_\zeta|}$), there is a function

$$c_\zeta : X_\zeta \rightarrow \kappa$$

such that

$$c_\zeta(x) \notin F_\zeta(x) \text{ for } x \in X_\zeta, \text{ and } |I_{c_\zeta}(A_\alpha)| = \kappa \text{ for all } A_\alpha \in \mathcal{A}_\zeta.$$

If $A_\alpha \in \mathcal{A}_\zeta$, then $|A_\alpha \cap X_{<\zeta}| < \kappa$ by (M2), and so $|I_{c_\zeta}(A_\alpha)| = \kappa$ implies

$$|I_{c_{\leq \zeta}}(A_\alpha)| = \kappa.$$

Since $\mathcal{I}$ is a principal ideal, we can pick $G_\zeta(A_\alpha) \in \mathcal{I} \cap [I_{c_{\leq \zeta}}(A_\alpha)]^\kappa$.

Finally take $c = \cup\{c_\zeta : \zeta < \text{cf}(\lambda)\}$.

Then c witnesses ($\bullet_\lambda$). ($\mathcal{I}1$) is clear from the construction. As for ($\mathcal{I}2$), assume that $A_\alpha \in \mathcal{A}_\zeta$. Then $G_\zeta(A_\alpha) \subset I_{c_{\leq \zeta}}(A_\alpha)$ by (6), and if $x \in A_\alpha \setminus X_{\leq \zeta}$, then $c(x) \notin G_\zeta(A_\alpha)$. So $G_\zeta(A_\alpha) \subset I_c(A)$ as well.

So we completed the proof of Theorem 4.1. □

REFERENCES

- [1] P. Erdős, F. Galvin and A. Hajnal: *On set-systems having large chromatic number and not containing prescribed subsystems*, Infinite and finite sets (Colloq., Keszthely, 1973; dedicated to P. Erdős on his 60th birthday), Vol. I; Colloq. Math. Soc. János Bolyai, Vol. 10, pp. 425–513, North-Holland, Amsterdam, 1975
- [2] P. Erdős and A. Hajnal: *On a property of families of sets*, Acta Math. Acad. Sci. Hung., 12,(1961) 87-124
- [3] P. Erdős and A. Hajnal: *On the chromatic number of graphs and set-systems*, Acta Math. Acad. Sci. Hung., 17,(1966) 159-229
- [4] P. Erdős, A. Hajnal and B. Rothchild: *On chromatic numbers of graphs and set-systems*, Proceedings of the 1971 Cambridge Summer School, Springer Lecture Notes in Math., no. 337, (1971), pp. 531-538.
- [5] A. Hajnal, I. Juhász, L. Soukup and Z. Szentmiklóssy: *Conflict free colorings of (strongly) almost disjoint set-systems* Acta Math. Hungar. 131(2011), Pages 230-274
- [6] P. Komjáth: *Dense systems of almost-disjoint sets*. Finite and infinite sets, (Eger, 1981), 527–536, Coll Math Soc. J. Bolyai, 10, 1984.
- [7] P. Komjáth: *Families close to disjoint ones*, Acta Math. Hungar. 43 (1984), pp 199–207.
- [8] P. Komjáth: *Conflict free colorings of nonuniform systems of infinite sets*, Acta Math. Hungar. 133(2011), 4, pp 305-310.
- [9] K. Kunen: *Set Theory*, North-Holland, New York, 1980.
- [10] E. W. Miller: *On a property of families of sets*, Comptes Rendus Varsovie, 30, (1937), pp 31-38.
- [11] S. Shelah: *A compactness theorem in singular cardinals, free algebras, Whitehead problem and transversals*, Israel Journal of Mathematics, 21 (1975), pp 319–349.
- [12] S. Shelah: *The Generalized Continuum Hypothesis revisited*, Israel Journal of Mathematics, 116 (1998) pp 285-321.
- [13] P. J. Szeptycki: *Transversals for strongly almost disjoint families*, Proc. Amer. Math. Soc., 135 (2007), 2273–2282.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES
E-mail address: `soukup@renyi.hu`
URL: `http://www.renyi.hu/~soukup`